Complete Summary

TITLE

Chronic stable coronary artery disease (CAD): percentage of patients who were screened for diabetes.

SOURCE(S)

American College of Cardiology, American Heart Association, Physician Consortium for Performance Improvement[™]. Clinical performance measures: chronic stable coronary artery disease. Tools developed by physicians for physicians. Chicago (IL): American Medical Association (AMA); 2005. 8 p. [18 references]

Measure Domain

PRIMARY MEASURE DOMAIN

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the Measure Validity page.

SECONDARY MEASURE DOMAIN

Does not apply to this measure

Brief Abstract

DESCRIPTION

This measure is used to assess the percentage of patients with chronic stable coronary artery disease (CAD) who were screened for diabetes.

RATIONALE

According to American Diabetes Association (ADA), American College of Endocrinology (ACE), and American Association of Clinical Endocrinologists (AACE) guidelines, screening for diabetes is recommended in patients who are considered high risk (e.g., coronary artery disease [CAD]).

PRIMARY CLINICAL COMPONENT

Coronary artery disease (CAD); screening for diabetes

DENOMINATOR DESCRIPTION

All patients with coronary artery disease (CAD) who do not have documented diabetes

NUMERATOR DESCRIPTION

Patients from the denominator who were screened for diabetes

Evidence Supporting the Measure

EVIDENCE SUPPORTING THE CRITERION OF QUALITY

- A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence
- One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

NATIONAL GUIDELINE CLEARINGHOUSE LINK

 ACC/AHA 2002 guideline update for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1999 Guidelines for the Management of Patients With Chronic Stable Angina).

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Variation in quality for the performance measured

EVIDENCE SUPPORTING NEED FOR THE MEASURE

American College of Endocrinology Consensus Statement on guidelines for glycemic control. Endocr Pract2002;8(Suppl 1):6-11.

American Diabetes Association. Clinical practice recommendations 2003. Screening for type 2 diabetes (position statement). Diabetes Care2003; 26(suppl 1): 21-4.

Communication from American Association of Clinical Endocrinologists. January 2003.

State of Use of the Measure

STATE OF USE

Pilot testing

CURRENT USE

Internal quality improvement

Application of Measure in its Current Use

CARE SETTING

Ambulatory Care Community Health Care Managed Care Plans Physician Group Practices/Clinics Rural Health Care

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Advanced Practice Nurses Physician Assistants Physicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Individual Clinicians

TARGET POPULATION AGE

Patients of all ages with the diagnosis of chronic stable coronary artery disease (CAD)

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

- Approximately 13 million Americans are living with coronary artery disease (CAD).
- More than 1 million Americans had a new or recurrent coronary attack in 2001.

EVIDENCE FOR INCIDENCE/PREVALENCE

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

- Chronic stable coronary artery disease (CAD) is the leading cause of mortality in the United States, accounting for almost 1 in 5 deaths.
- For individuals with CAD, the risk of another heart attack, stroke, and other serious complications is substantial.

EVIDENCE FOR BURDEN OF ILLNESS

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

UTILIZATION

Within the past 2 decades, the number of short-stay hospital discharges for individuals with coronary artery disease (CAD) increased by almost 18%.

EVIDENCE FOR UTILIZATION

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

COSTS

The total annual cost of coronary artery disease (CAD) in the United States is approximately \$130 billion.

EVIDENCE FOR COSTS

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Living with Illness

IOM DOMAIN

Effectiveness

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

All patients with coronary artery disease (CAD) who do not have documented diabetes

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

All patients with coronary artery disease (CAD) who do not have documented diabetes

Exclusions

Patients with documented diabetes

RELATIONSHIP OF DENOMINATOR TO NUMERATOR

All cases in the denominator are equally eligible to appear in the numerator

DENOMINATOR (INDEX) EVENT

Clinical Condition

DENOMINATOR TIME WINDOW

Time window follows index event

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

Patients from the denominator who were screened for diabetes*

Exclusions

None

MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

^{*}Screening for diabetes is usually done by fasting blood glucose or 2-hour glucose tolerance testing. Clinical recommendations indicate screening should be considered at 3-year intervals.

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

NUMERATOR TIME WINDOW

Fixed time period

DATA SOURCE

Medical record

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

None

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

Unspecified

Identifying Information

ORIGINAL TITLE

Screening for diabetes.

MEASURE COLLECTION

The Physician Consortium for Performance Improvement Measurement Sets

MEASURE SET NAME

American College of Cardiology, American Heart Association, and Physician Consortium for Performance Improvement: Chronic Stable Coronary Artery Disease Physician Performance Measurement Set

SUBMITTER

American Medical Association on behalf of the American College of Cardiology, the American Heart Association, and the Physician Consortium for Performance Improvement

DEVELOPER

American College of Cardiology - Medical Specialty Society American Heart Association Physician Consortium for Performance Improvement

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2003 Jan

REVISION DATE

2005 Aug

MEASURE STATUS

This is the current release of the measure.

This measure updates a previous version: American College of Cardiology, American Heart Association, Physician Consortium for Performance Improvement. Clinical performance measures. Chronic stable coronary artery disease. Chicago (IL): American Medical Association (AMA); 2003. 8 p.

SOURCE(S)

American College of Cardiology, American Heart Association, Physician Consortium for Performance Improvement™. Clinical performance measures: chronic stable

coronary artery disease. Tools developed by physicians for physicians. Chicago (IL): American Medical Association (AMA); 2005. 8 p. [18 references]

MEASURE AVAILABILITY

The individual measure, "Screening for Diabetes," is published in the "Clinical Performance Measures: Chronic Stable Coronary Artery Disease." This document and technical specifications are available in Portable Document Format (PDF) from the American Medical Association (AMA)-convened Physician Consortium for Performance Improvement Web site: www.physicianconsortium.org.

For further information, please contact AMA staff by e-mail at cgi@ama-assn.org.

COMPANION DOCUMENTS

The following are available:

- Physician Consortium for Performance Improvement. Introduction to physician performance measurement sets. Tools developed by physicians for physicians. Chicago (IL): American Medical Association (AMA); 2001 Oct. 21 p. This document is available from the American Medical Association (AMA) Clinical Quality Improvement Web site: www.ama-assn.org/go/guality.
- Physician Consortium for Performance Improvement. Principles for performance measurement in health care. A consensus statement. [online]. Chicago (IL): American Medical Association (AMA), Joint Commission on the Accreditation of Healthcare Organizations (JCAHO), National Committee for Quality Assurance (NCQA); [3 p]. This document is available from the AMA Clinical Quality Improvement Web site: www.ama-assn.org/go/quality.
- Physician Consortium for Performance Improvement. Desirable attributes of performance measures. A consensus statement. [online]. American Medical Association (AMA), Joint Commission on Accreditation of Healthcare Organizations (JCAHO), National Committee for Quality Assurance (NCQA); 1999 Apr 19 [cited 2002 Jun 19]. [5 p]. This document is available from the AMA Clinical Quality Improvement Web site: www.ama-assn.org/qo/quality.

For further information, please contact AMA staff by e-mail at cgi@ama-assn.org.

NQMC STATUS

This NQMC summary was completed by ECRI on September 26, 2003. The information was verified by the measure developer on January 28, 2004. This NQMC summary was updated by ECRI on September 28, 2005.

COPYRIGHT STATEMENT

Physician Performance Measures (Measures) and related data specifications, developed by the Physician Consortium for Performance Improvement (the Consortium), are intended to facilitate quality improvement activities by physicians.

These Measures are intended to assist physicians in enhancing quality of care. Measures are designed for use by any physician who manages the care of a patient for a specific condition or for prevention. These performance Measures are not clinical guidelines and do not establish a standard of medical care. The Consortium has not tested its Measures for all potential applications. The Consortium encourages the testing and evaluation of its Measures.

Measures are subject to review and may be revised or rescinded at any time by the Consortium. The Measures may not be altered without the prior written approval of the Consortium. Measures developed by the Consortium, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g., use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the Measures require a license agreement between the user and American Medical Association, on behalf of the Consortium. Neither the Consortium nor its members shall be responsible for any use of these Measures.

THE MEASURES ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.

© 2005 American Medical Association. All Rights Reserved.

© 2006 National Quality Measures Clearinghouse

Date Modified: 9/25/2006



